arXiv:1701.02237v1 [math.MG] 4 Jan 2017

COMPARING VOLUMES BY CONCURRENT CROSS-SECTIONS OF COMPLEX LINES: A BUSEMANN-PETTY TYPE PROBLEM

ERIC L. GRINBERG
UNIVERSITY OF MASSACHUSETTS, BOSTON
BOSTON MA 02125, USA

Abstract. We consider the problem of comparing the volumes of two star bodies in an even-dimensional euclidean space $\mathbb{R}^{2n} = \mathbb{C}^n$ by comparing their cross sectional areas along complex lines (special 2-dimensional real planes) through the origin. Under mild symmetry conditions on one of the bodies a Busemann-Petty type theorem holds. Quaternionic and Octonionic analogs also hold. The argument relies on integration in polar coordinates coupled with Jensen's inequality. Along the way we provide a criterion that detects which centered bodies are *circular*. i.e., stabilized by multiplication by complex numbers of unit modulus. Our goal is to present a Busemann-Petty type result with a minimum of required background and, in addition, to suggest characterizations of classes of star bodies by means of integral geometric inequalities.

Domains with mild symmetry

Definition. *Let D be domain in $\mathbb{C}^n$. Then D is* **a circular domain, with center** $\longrightarrow 0$ *if and only if it is stabilized by multiplication by complex numbers of unit modulus, i.e.,*

$$z \in D,\, \theta \in \mathbb{R} \quad \Longrightarrow \quad e^{\sqrt{-1}\theta} z \in D.$$

D is said to be circular with center $\overrightarrow{c}$ when $D - \overrightarrow{c}$ is circular with center $\overrightarrow{0}$.

Thus the origin centered unit ball $\{\vec{z} \in \mathbb{C}^n \,\Big|\, \|\vec{z}\| = 1\}$ and the standard unit polydisc,

$$P \equiv \{\vec{z} \in \mathbb{C}^n \,\Big|\, |z_i| \leq 1, i = 1 \dots n\},$$

are both circular.

In the convex geometry literature circular bodies are sometimes said to have R_θ symmetry. In the functional analysis literature one sometimes speaks of a *balanced set*.

Theorem 1. *Let D be a star-shaped origin symmetric domain in $\mathbb{C}^n$ and let $G_{1,n}$ be the set of 1-dimensional vector subspaces of $\mathbb{C}^n$, that is, the set of complex lines through the origin in $\mathbb{C}^n$. Then the volume of D dominates the mean of the n^{th} power of cross-sectional areas of D by lines in $G_{1,n}$, with equality precisely when D is circular:*

$$vol(D) \geq \int_{\ell \in G_{1,n}} (area(D \cap \ell))^n \, d\ell.$$

This suggests the quest of characterizing all 'classes' of bodies by geometric integral inequalities and identities, where *classes*, *geometric*, and *integral* are left to be made more precise (not to mention *all*). For instance, characterizations of centered bodies in $\mathbb{R}^n$ are found in [Gardner] and [Schneider]. We seek a mapping (functor?) from classes of bodies to geometric inequalities, perhaps in the spirit of [Zalc], where a parallel is drawn between mean value properties and differential equations. For some results in this direction see [Dann-Zy].

Proof. Viewing the ambient space $\mathbb{C}^n$ as $\mathbb{R}^{2n}$, we can compute the volume of the region D in polar coordinates. We recall that the *radial function* of D, $\rho_D(\omega D)$, is defined by

$$\rho_D(\omega) = \max\{t \in [0, +\infty) \,|\, t\omega \in D\}.$$

With this notation we have

$$\mathrm{vol}(D) = \int_{\omega \in S^{2n-1}} \left(\rho_D(\omega)\right)^{2n} \, d\omega,$$

where $d\omega$ is the suitably normalized rotation invariant measure on the sphere S^{2n-1}. Since multiplication by a phase factor stabilizes the sphere and preserves its measure,

$$\int_{\omega \in S^{2n-1}} f(\omega)\, d\omega = \int_{\omega \in S^{2n-1}} f(e^{i\theta}\omega)\, d\omega$$

for any θ. So

$$\begin{aligned}
\mathrm{vol}(D) &= \textstyle\int_{\omega \in S^{2n-1}} \left(\rho_D(\omega)\right)^{2n} \, d\omega \\
&= \textstyle\int_{\omega \in S^{2n-1}} \left(\rho_D(e^{i\theta}\omega)\right)^{2n} \, d\omega \\
&= \textstyle\int_{\omega \in S^{2n-1}} \int_0^{2\pi} \left(\rho_D(e^{i\theta}\omega)\right)^{2n} \left(\frac{1}{2\pi}\right) d\theta \, d\omega \\
&= \textstyle\int_{\omega \in S^{2n-1}} \int_0^{2\pi} \left\{\left(\rho_D(e^{i\theta}\omega)\right)^{2}\right\}^{n} \left(\frac{1}{2\pi}\right) d\theta \, d\omega \\
&\geq \textstyle\int_{\omega \in S^{2n-1}} \left\{\int_0^{2\pi} \left(\rho_D(e^{i\theta}\omega)\right)^{2} \frac{1}{2\pi} d\theta\right\}^{n} \, d\omega \\
&= \textstyle\int_{\omega \in S^{2n-1}} \{area(D \cap \ell(\omega))\}^{n} \, d\omega \,.
\end{aligned}$$

The next to last line above is obtained by applying Jensen's inequality [Rudin] to the convex function $x \mapsto x^n$ on $[0, +\infty)$. Notice that equality holds iff $\rho_D(e^{i\theta}\omega) = \rho_D(\omega)$ for almost all (and hence for all) θ and ω (by continuity), i.e., if and only if D is a circular region. □

Theorem 2. *Let A and B be origin-symmetric star bodies in $\mathbb{C}^n$ with A circular. If for every complex line ℓ through the origin in $\mathbb{C}^n$ we have*

$$area(A \cap \ell) \,\leq\, area(B \cap \ell),$$

then A has smaller volume than B.

This is a variation on the classical *Busemann-Petty* problem. It is powered entirely by integration over polar coordinates (adapted to complex geometry) together with Jensen's inequality.

Proof. By the previous theorem (statement and proof), the volume of A is equal to the n^{th} power mean of its concurrent complex line cross sectional areas, which is dominated by the corresponding mean of cross sections of B, which is dominated by the volume of B:

$$\begin{aligned} \mathrm{vol}(A) &= \int_{\ell \in G_{1,n}} (\mathrm{area}(A \cap \ell))^n \, d\ell \\ &\leq \int_{\ell \in G_{1,n}} (\mathrm{area}(B \cap \ell))^n \, d\ell \\ &\leq \mathrm{vol}(B). \end{aligned}$$

□

Note: a similar argument allows comparison of bodies K, L in $\mathbb{R}^{4n}$ by cross-sections along quaterionic lines (assuming K is quaternion-circular, or S^3-circular.)

Acknowledgement

The author wishes to thank Susanna Dann, David Feldman, Daniel Klain, Mehmet Orhon and others for helpful discussions.

References

[BFM] F. Barthe; M. Fradelizi; B. Maurey, A short solution to the Busemann-Petty problem, Positivity **3** (1999), 95–100.

[BP] H. Busemann, C. M. Petty, Problems on convex bodies, Math. Scand., **4** (1956), 88–94.

[Dann] S. Dann, The Busemann-Petty problem in the complex hyperbolic space, Math. Proc. Cambridge Philos. Soc. **155** (2013), no. 1, 155–172

[Dann-Zy] Dann, Susanna; Zymonopoulou, Marisa Sections of convex bodies with symmetries. Adv. Math. 271 (2015), 112–152.

[Gardner] R.J. Gardner, *Geometric Tomography*, second edition, Cambridge University Press, 2006

[Gr] E. Grinberg, Isoperimetric inequalities and identities for k-dimensional cross-sections of convex bodies, Math. Ann. **291** (1991), no. 1, 75-86.

[KKZ] A. Koldobsky, H. König, M. Zymonopoulou, The complex Busemann-Petty problem on sections of convex bodies, Advances in Mathematics, **218** (2008), 352–367.

[Rubin1] B. Rubin, The lower dimensional Busemann-Petty problem for bodies with the generalized axial symmetry, Israel J. Math. **173** (2009), 213-233.

[Rubin2] B. Rubin, Comparison of volumes of convex bodies in real, complex, and quaternionic spaces. Adv. Math. **225** (2010), no. 3, 1461-1498.

[Rudin] W. Rudin, Real and Complex Analysis, 3rd Ed., Chp. 3, McGraw-Hill (1986).

[Schneider] Schneider, R., *Convex Bodies The BrunnMinkowski Theory.* 2nd ed., Cambridge University Press, Cambridge, 2014.

[Zalc] L. Zalcman, Mean values and differential equations, Israel J. Math. **14** (1973), 339-352.

[Zy] M. Zymonopoulou, The modified complex Busemann-Petty problem on sections of convex bodies, Positivity **13** (2009), no. 4, 717–733.